



**HAL**  
open science

## Understanding the History of the Domestic Cycle of *Toxoplasma gondii* in South America

Lokman Galal, Frédéric Arieu, Meriadeg Ar Gouilh, Marie-Laure Dardé, Azra Hamidović, Franck Letourneur, Franck Prugnolle, Aurélien Mercier

► **To cite this version:**

Lokman Galal, Frédéric Arieu, Meriadeg Ar Gouilh, Marie-Laure Dardé, Azra Hamidović, et al.. Understanding the History of the Domestic Cycle of *Toxoplasma gondii* in South America. 3rd Environmental Toxoplasmosis Workshop, May 2024, BERLIN, Germany. hal-04598103

**HAL Id: hal-04598103**

**<https://unilim.hal.science/hal-04598103>**

Submitted on 3 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Understanding the History of the Domestic Cycle of *Toxoplasma gondii* in South America

Lokman Galal <sup>1</sup>, Frédéric Arieu <sup>2</sup>, Meriadeg Ar Gouilh <sup>3,4,5</sup>, Marie-Laure Dardé <sup>1,6</sup>, Azra Hamidović <sup>1</sup>, Franck Letourneur <sup>7</sup>, Franck Prugnolle <sup>8</sup> and Aurélien Mercier <sup>1,6</sup>

<sup>1</sup>InsermU1094, IRD U270, Univ. Limoges, CHU Limoges, EpiMaCT—Epidemiology of chronic diseases in tropical zone, Institute of Epidemiology and Tropical Neurology, OmegaHealth, Limoges, France.

<sup>2</sup>Université de Paris, Institut Cochin, Inserm U1016, Service de Parasitologie Hôpital Cochin, 75014 Paris, France.

<sup>3</sup>DYNAMICURE U1311 INSERM, Université de Caen Normandie, UNICAEN, UNIROUEN, 14000 Caen, France.

<sup>4</sup>Laboratoire de Virologie, Centre Hospitalo-Universitaire, Avenue Georges Clémenceau, 14000 Caen, France.

<sup>5</sup>Institut de Recherche pour le Développement (IRD), Maladies Infectieuses et vecteurs : Ecologie, Génétique, Evolution et Contrôle (MIVEGEC) (Université de Montpellier—IRD 224—CNRS 5290), Montpellier, France.

<sup>6</sup>Centre National de Référence (CNR) Toxoplasmose/Toxoplasma Biological Center (BRC), Centre Hospitalier-Universitaire Dupuytren, Limoges, France.

<sup>7</sup>Plate-Forme Séquençage et Génomique, Institut Cochin, Inserm U1016, Université de Paris, 22 rue Méchain, 75014 Paris, France.

<sup>8</sup>IRL REHABS, International Research Laboratory REHABS, CNRS-NMU-UCBL, Nelson Mandela University George Campus, George 6531, South Africa.

**e-mail:** [lokman.galal@unilim.fr](mailto:lokman.galal@unilim.fr)

---

A growing body of evidence supports the significant role of *Toxoplasma gondii* genotype in determining disease severity. However, isolating and characterizing *T. gondii* strains infecting humans remains challenging. Clinical severity of toxoplasmosis varies geographically, with South America experiencing the highest burden, coinciding with the continent's remarkable *T. gondii* genetic diversity. In our study, we analyzed 156 *T. gondii* genomes and we provide the first direct estimates of *T. gondii*'s mutation rate and generation time. Our findings offer insights into the epidemiology, ecology, and evolutionary history of South American *T. gondii* populations. We identified three distinct categories: (1) wild populations, transmitted through the sylvatic cycle involving wild felids and likely poorly adapted to domestic cats. This includes the "Amazonian" population, linked to various clinical forms in immunocompetent individuals, and the "Pan-American" population, isolated from

wildlife and not yet associated with severe disease forms, (2) domestic intercontinental clonal lineages (types I, II, III, and Africa 1), introduced to the continent with the domestic cat and house mouse by European sailors. (3) hybrids of the first two categories, probably well adapted to domestic cats, and representing the majority of strains found in South America's domestic environment. We discuss how hybridization events and strong selection for a unique haplotype comprising only 0.16% of the *T. gondii* genome on chromosome 1a are likely driving the public health challenge associated with toxoplasmosis in South America.

**Galal, L., Ariey, F., Gouilh, M. A., Dardé, M. L., Hamidović, A., Letourneur, F., ... & Mercier, A. (2022). A unique *Toxoplasma gondii* haplotype accompanied the global expansion of cats. *Nature Communications*, 13(1), 5778.**